




TMDL for Sediment in the Non-Tidal Marsh Run

What You Need to Know

Background

The [Total Maximum Daily Load, or TMDL, for sediment in the Marsh Run watershed](#) establishes an annual load limit for total suspended solids (TSS) to the non-tidal streams within the watershed.



The [Marsh Run watershed](#), as defined by the State of Maryland, is located entirely within Washington County, MD. Washington County is bordered by Mason and Dixon’s line and Pennsylvania to the north and by the far shore of the Potomac River, Virginia, and West Virginia to the south. It extends eastward to South Mountain and Frederick County and westward to Sideling Hill Creek and Allegany County. The watershed is located within the Highland eco-region.



Impaired aquatic life and wildlife in the watershed was first identified in 2004 based on results from the Maryland Biological Stream Survey (MBSS), a randomized survey of stream health. As part of the MBSS, streams are scored against reference watersheds where habitat and aquatic diversity is high, using two biological indices: the Benthic Index of Biotic Integrity (BIBI) which looks at the biological community in the bottom sediments, and the Fish Index of Biotic Integrity (FIBI).

Watershed ID	Maryland 8-Digit: 02140503
Watershed size	13,430 acres not including wetlands/water
Waterbody type	1 st - through 4 th -order non-tidal streams
Waterbody designated use not being met	Aquatic life and wildlife
Reason for impairment	Stream biology impacted by excessive sediment
TMDL Baseline year	2009
Overall sediment reduction percent	31%

A biological stressor identification (BSID) analysis was conducted in 2015 to identify possible causes of the stream degradation. Using MBSS data, [the BSID](#) showed sediment indicators, instream habitat, and water chemistry pollutants as potential causes. Based on this assessment, the Marsh Run watershed was listed as impaired for sediment, as well as sulfates on [Maryland’s 2016 Integrated Report of Surface Water Quality](#). The [non-tidal sediment TMDL](#) was approved on September 12, 2019.

TMDL

The TMDL for sediment in the Marsh Run watershed, was established at a level to ensure acceptable biological integrity in the watershed’s streams.

The TMDL was developed with a reference watershed approach using loading results from the Phase 5.3.2 Chesapeake Bay Watershed Model. Annual sediment loads

from reference watersheds—those with good biological integrity—were compared to predicted loads under a modeled all forest scenario to establish an acceptable ratio of current loadings to loadings in a natural condition. This ratio, known as the forest normalized load (FNL), was also calculated for the Marsh Run watershed, and the TMDL was established based on the reduction needed to achieve the reference FNL.

Allocations

Allocations to point sources such as wastewater treatment plants and regulated stormwater, are called Wasteload Allocations (WLAs), and allocations to nonpoint sources, like cropland, are called Load Allocations (LAs). Sector load reductions in this TMDL were assigned using the controllable load methodology from Maryland's Phase II Watershed Implementation Plan (WIP) for the Chesapeake Bay. This methodology assigns reductions to controllable loads (e.g., agriculture & urban) and gives credit to existing implementation efforts, resulting in different percent reductions for different source categories.

The watershed has no municipal dischargers with a permit regulating the discharge of sediment. Regulated stormwater

sources include Phase II municipal separate storm sewer system (MS4) and several entities covered under general permits. The WLAs are described in detail in the TMDL’s [technical memorandum on point sources](#). Regulated stormwater WLAs represent less than 1% of the total TMDL.

The LAs for this TMDL, as presented in the [technical memorandum on nonpoint sources](#), account for the remaining portion of the TMDL and are assigned to unregulated urban, agricultural, and natural source categories.

The Baseline and TMDL equations for the Marsh Run watershed, including source categories and allocations are provided in the equation below.

Baseline Equation:	10,817	=	449	+	8,894	+	130	+	1,344	+	0	TSS tons / year
	Baseline Load		Forest		Agriculture		Unregulated Urban		Stormwater		Wastewater	
TMDL Equation:	7,418	=	449	+	6,366	+	54	+	548	+	0	TSS tons / year
	TMDL		Forest LA		Agriculture LA		Unregulated Urban LA		Stormwater WLA		Wastewater WLA	
Reduction from baseline	31%		0%		28%		58%		59%		0%	

Note: The loadings in this TMDL are expressed as Edge-of-Stream, or EOS, loads, based on the Phase 5.3.2 Chesapeake Bay Watershed Model

Next Steps

Most of the sediment reductions in this TMDL are assigned to agricultural and urban stormwater sources. Implementation of these reductions will occur in parallel with efforts to fully implement the 2010 Chesapeake Bay TMDLs by 2025. While the endpoints of the TMDLs are different—tidal water quality actions will result in progress toward both goals.

Sediment reductions from agricultural sources are usually achieved by managing runoff and erosion with best management practices (BMPs) such as conservation tillage, and riparian buffers. Statewide programs and regulations that promote agricultural BMPs will reduce sediment loads in the Marsh Run watershed. The Chesapeake and Atlantic Coastal Bays Trust Fund, for example, makes funds available for planting cover crops.

For urban stormwater, sediment reductions are typically achieved by addressing water quality and quantity with stormwater BMP retrofits. Retrofits include the modification of existing stormwater ponds, the installation of new structural BMPs, tree planting and stream restoration. Individually

permitted MS4 jurisdictions are required to develop plans for implementing the sediment reductions from this TMDL. The BMPs described in these plans may also be used for meeting permit impervious area restoration requirements.

While this TMDL establishes a sediment loading target for the watershed, and sediment load reductions are an important tool for tracking progress, the measure of its successful implementation will be its effect on in-stream biological health. The watershed cannot be classified as meeting water quality standards until it is demonstrated that aquatic life is no longer impaired by sediment.